


## 4.2.1 Alcohols

The alcohols have relatively low volatility due to their ability to form hydrogen bond between alcohol molecules.

The smaller alcohols (up to 3 carbons) are soluble in water because they can form hydrogen bonds with water. The longer the hydrocarbon chain the less soluble the alcohol.


General formula alcohols  $C_nH_{2n+1}OH$




### Uses of alcohols

Ethanol is 'alcohol' in alcoholic drinks. Ethanol is commonly used as a solvent in the form of methylated spirits. Methanol is used as a petrol additive to improve combustion and is increasing important as a feedstock in the production of organic chemicals;


### Classifying Alcohols



Propan-1-ol  
Primary



Propan-2-ol  
Secondary



methylpropan-2-ol  
Tertiary

Primary alcohols are alcohols where 1 carbon is attached to the carbon adjoining the oxygen

Secondary alcohols are alcohols where 2 carbon are attached to the carbon adjoining the oxygen


Tertiary alcohols are alcohols where 3 carbon are attached to the carbon adjoining the oxygen

### Reactions of alcohols

#### Complete Combustion

In excess oxygen alcohols will burn with complete combustion

The products of *complete* combustion are  $CO_2$  and  $H_2O$ .



## Oxidation reactions of the alcohols

Potassium dichromate  $K_2Cr_2O_7$  is an oxidising agent that causes alcohols to oxidise.


The exact reaction, however, depends on the type of alcohol, i.e. whether it is primary, secondary, or tertiary, and on the conditions.

### Partial Oxidation of Primary Alcohols

**Reaction:** primary alcohol  $\rightarrow$  aldehyde


**Reagent:** potassium dichromate (VI) solution and dilute sulphuric acid.

**Conditions:** (use a limited amount of dichromate) warm gently and **distil** out the aldehyde as it forms:



An aldehyde's name ends in **-al**

It always has the C=O bond on the first carbon of the chain so it does not need an extra number



Observation: the orange dichromate ion ( $Cr_2O_7^{2-}$ ) reduces to the green  $Cr^{3+}$  ion

Write the oxidation equations in a simplified form using [O] which represents O from the oxidising agent

When writing the formulae of aldehydes in a condensed way write **CHO** and not COH e.g.  $CH_3CH_2CHO$


## Distillation

In general used as separation technique to separate an organic product from its reacting mixture. Need to collect the distillate of the approximate boiling point range of the desired liquid.

Note the bulb of the thermometer should be at the T junction connecting to the condenser to measure the correct boiling point

Note the water goes in the bottom of the condenser to go against gravity. This allows more efficient cooling and prevents back flow of water.

Electric heaters are often used to heat organic chemicals. This is because organic chemicals are normally highly flammable and could set on fire with a naked flame.





It's important to be able to draw and label this apparatus accurately. Don't draw lines between flask, adaptor and condenser.

## Full Oxidation of Primary Alcohols

**Reaction:** primary alcohol  $\rightarrow$  carboxylic acid

**Reagent:** potassium dichromate(VI) solution and dilute sulphuric acid

**Conditions:** use an excess of dichromate, and **heat under reflux**: (distill off product after the reaction has finished)




Observation: the orange dichromate ion ( $\text{Cr}_2\text{O}_7^{2-}$ ) reduces to the green  $\text{Cr}^{3+}$  ion

## Reflux

Reflux is used when heating organic reaction mixtures for long periods. The condenser prevents organic vapours from escaping by condensing them back to liquids.

**Never seal the end of the condenser** as the build up of gas pressure could cause the apparatus to explode. This is true of any apparatus where volatile liquids are heated including the distillation set up

Anti-bumping granules are added to the flask in both distillation and reflux to prevent vigorous, uneven boiling by **making small bubbles** form instead of large bubbles




## Oxidation of Secondary Alcohols

**Reaction:** secondary alcohol  $\rightarrow$  ketone

**Reagent:** potassium dichromate(VI) solution and dilute sulphuric acid.


**Conditions:** heat under reflux



Propanone

Ketones end in **-one**

When ketones have 5C's or more in a chain then it needs a number to show the position of the double bond. E.g. pentan-2-one



Observation: the orange dichromate ion ( $\text{Cr}_2\text{O}_7^{2-}$ ) reduces to the green  $\text{Cr}^{3+}$  ion

There is no further oxidation of the ketone under these conditions.

**Tertiary alcohols** cannot be oxidised at all by potassium dichromate: This is because there is no hydrogen atom bonded to the carbon with the OH group

## Reaction of alcohols with Dehydrating agents

**Reaction:** Alcohol  $\rightarrow$  Alkene


**Reagents:** Concentrated Sulphuric or Phosphoric acids

**Conditions:** warm (under reflux)


**Role of reagent:** dehydrating agent/catalyst

**Type of reaction:** acid catalysed elimination

Dehydration Reaction: removal of a water molecule from a molecule



Some 2° and 3° alcohols can give more than one product, when the double bond forms between different carbon atoms



## Substitution reactions of Alcohols to form Haloalkanes

A mixture of a halide ions with concentrated acid NaCl + H<sub>2</sub>SO<sub>4</sub> can be used for substituting a halogen on to an alcohol

Various other halogenating compounds can be used to substitute the –OH group for a halogen

**Reaction:** Alcohol → Haloalkane

**Reagents:** Concentrated Sulphuric and sodium halide

