



## 18. Carbonyls: Aldehydes and Ketones

Carbonyls are compounds with a C=O bond. They can be either aldehydes or ketones



If the C=O is on the end of the chain with an H attached it is an aldehyde.  
The name will end in **-al**

CH<sub>3</sub>CHO ethanal



If the C=O is in the middle of the chain it is a ketone.  
The name will end in **-one**


CH<sub>3</sub>COCH<sub>3</sub> propanone

### Intermolecular forces in Carbonyls

Pure carbonyls cannot hydrogen bond, but bond instead by **permanent dipole forces**.


### Solubility in water

The smaller carbonyls are soluble in water because they can form hydrogen bonds with water.



### Reactions of carbonyls

The C=O bond is polarised because O is more electronegative than carbon. The positive carbon atom attracts nucleophiles.



In comparison to the C=C bond in alkenes, the C=O is stronger and does not undergo addition reactions easily.

This is in contrast to the electrophiles that are attracted to the C=C.

### Oxidation Reactions


Potassium dichromate K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> is an oxidising agent that causes alcohols and aldehydes to oxidise.

Primary alcohol → aldehydes → carboxylic acid  
Secondary alcohol → ketones  
Tertiary alcohols do not oxidise


**Key point:** Aldehydes can be oxidised to carboxylic acids, but ketones cannot be oxidised.

### Oxidation of Aldehydes

**Reaction:** aldehyde → carboxylic acid  
**Reagent:** potassium dichromate (VI) solution and dilute sulphuric acid.  
**Conditions:** heat under reflux



### Full Equation for oxidation



Observation: the orange dichromate ion (Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup>) reduces to the green Cr<sup>3+</sup> ion

Aldehydes can also be oxidised using Fehling's solution or Tollen's Reagent. These are used as tests for the presence of aldehyde groups


### Tollen's Reagent

**Reagent:** Tollen's Reagent formed by mixing aqueous ammonia and silver nitrate. The active substance is the complex ion of [Ag(NH<sub>3</sub>)<sub>2</sub>]<sup>+</sup>.

**Conditions:** heat gently

**Reaction:** aldehydes only are oxidised by Tollen's reagent into a carboxylic acid and the silver(I) ions are reduced to silver atoms

**Observation:** with aldehydes, a silver mirror forms coating the inside of the test tube. Ketones result in no change.




### Fehling's solution

**Reagent:** Fehling's Solution containing blue Cu<sup>2+</sup> ions.

**Conditions:** heat gently

**Reaction:** aldehydes only are oxidised by Fehling's Solution into a carboxylic acid and the copper ions are reduced to copper(I) oxide.

**Observation:** Aldehydes: Blue Cu<sup>2+</sup> ions in solution change to a red precipitate of Cu<sub>2</sub>O. Ketones do not react.



## Reduction of carbonyls

**Reagents:** LiAlH<sub>4</sub> in dry ether


**Conditions:** Room temperature and pressure

**Type of reaction:** Reduction


**Role of reagent:** Reducing agent

Reducing agents such as NaBH<sub>4</sub> (sodium tetrahydridoborate) or LiAlH<sub>4</sub> (lithium tetrahydridoaluminate) will reduce carbonyls to alcohols.

Aldehydes will be reduced to primary alcohols



Ketones will be reduced to secondary alcohols.



## Addition of hydrogen cyanide to carbonyls to form hydroxynitriles



**Reaction:** carbonyl → hydroxynitrile

**Reagent:** HCN in presence of KCN



**Conditions:** Room temperature and pressure

**Mechanism:** nucleophilic addition


When naming hydroxy nitriles the CN becomes part of the main chain



2-hydroxy-2-methylpropanenitrile




2-hydroxypropanenitrile



The extra KCN increases the concentration of the CN<sup>-</sup> ion nucleophile needed for the first step of the mechanism

## Nucleophilic Addition Mechanism




## Reaction of carbonyls with iodine in presence of alkali

**Reagents:** Iodine and sodium hydroxide



**Conditions:** warm very gently

The product CHI<sub>3</sub> is a yellow crystalline precipitate with an antiseptic smell

Only carbonyls with a methyl group next to the C=O bond will do this reaction. Ethanal is the only aldehyde that reacts. More commonly is methyl ketones.



This reaction is called the Iodoform test



## Reaction with 2,4-dinitro phenylhydrazine

2,4-DNP reacts with both aldehydes and ketones. The product is an orange precipitate, It can be used as a test for a carbonyl group in a compound.

Use 2,4-DNP to identify if the compound is a carbonyl. Then to differentiate an aldehyde from a ketone use Tollen's reagent.

The melting point of the crystal formed can be used to help identify which carbonyl was used. Take the melting point of orange crystals product from 2,4-DNP. Compare melting point with known values in database

