

## 22. Analytical Techniques

### Chromatography

Chromatography is an analytical technique that separates components in a mixture between a mobile phase and a stationary phase.

Types of chromatography include:

- thin-layer chromatography (TLC) – a plate is coated with a solid and a solvent moves up the plate
- column chromatography (CC) – a column is packed with a solid and a solvent moves down the column
- gas chromatography (GC) – a column is packed with a solid or with a solid coated by a liquid, and a gas is passed through the column under pressure at high temperature.

Separation by column chromatography depends on the balance between solubility in the moving phase and retention in the stationary phase.

A solid stationary phase separates by adsorption,  
A liquid stationary phase separates by relative solubility

The mobile phase may be a liquid or a gas.  
The stationary phase may be a solid (as in thin-layer chromatography, TLC) or either a liquid or solid on a solid support (as in gas chromatography, GC)

In gas-liquid chromatography GC the **mobile** phase is an inert **gas** such as nitrogen, helium, argon.  
The **Stationary** phase is a **liquid** on an inert solid.

If the stationary phase was polar and the moving phase was non-polar e.g. Hexane. Then non-polar compounds would pass through the column more quickly than polar compounds as they would have a greater solubility in the non-polar moving phase.  
(Think about intermolecular forces)

Retention times and  $R_f$  values are used to identify different substances.

Method: **Thin-layer chromatography**

- Wearing gloves**, draw a **pencil line** 1 cm above the bottom of a TLC plate and mark spots for each sample, equally spaced along line.
- Use a capillary tube to add a **tiny drop** of each solution to a different spot and allow the plate to air dry.
- Add solvent to a chamber or large beaker with a lid so that is no more than **1cm in depth**
- Place the TLC plate into the chamber, **making sure that the level of the solvent is below the pencil line**. Replace the **lid to get a tight seal**.
- When the level of the solvent **reaches about 1 cm from the top of the plate**, remove the plate and mark the solvent level with a pencil. Allow the plate to **dry in the fume cupboard**.
- Place the plate under a **UV lamp** or if using amino acids spray with ninhydrin in order to see the spots. Draw around them lightly in pencil.
- Calculate the  $R_f$  values of the observed spots.

Wear plastic gloves to prevent contamination from the hands to the plate

**pencil line** – will not dissolve in the solvent

**tiny drop** – too big a drop will cause different spots to merge

**Depth** of solvent – if the solvent is too deep it will dissolve the sample spots from the plate


**lid** – to prevent evaporation of toxic solvent

Will get more accurate results if the solvent is allowed to rise to near the top of the plate but the  $R_f$  value can be calculated if the solvent front does not reach the top of the plate

dry in a **fume** cupboard as the solvent is toxic

**UV lamp** used if the spots are colourless and not visible


$R_f$  value =  $\frac{\text{distance moved by amino acid}}{\text{distance moved by the solvent}}$



## Column chromatography (CC)

### Simple column chromatography

- A glass tube is filled with the stationary phase usually silica or alumina in powder form to increase the surface area.
- A filter or plug is used to retain the solid in the tube. Solvent is added to cover all the powder.
- The mixture to be analysed is dissolved in a minimum of a solvent and added to the column.
- A solvent or mixture of solvents is then run through the column.
- The time for each component in the mixture to reach the end of the column is recorded (retention time)



HPLC stands for high performance liquid chromatography and is a type of column chromatography commonly used in industry.

HPLC: **stationary** phase is a **solid** silica

HPLC: **mobile** phase is a **liquid**

## Gas-Liquid Chromatography

Gas-liquid chromatography can be used to separate mixtures of volatile liquids.

The time taken for a particular compound to travel from the injection of the sample to where it leaves the column to the detector is known as its **retention time**. This can be used to identify a substance.


Some compounds have similar retention times so will not be distinguished.

Basic gas-liquid chromatography will tell us how many components there are in the mixture by the number of peaks. It will also tell us the abundance of each substance. The area under each peak will be proportional to the abundance of that component.

It is also possible for gas-liquid chromatography machine to be connected to a mass spectrometer, IR or NMR machine, enabling all the components in a mixture to be identified.

GC-MS is used in analysis, in forensics, environmental analysis, airport security and space probes.

In gas-liquid chromatography, the mobile phase is a gas such as helium and the stationary phase is a high boiling point liquid absorbed onto a solid.



Most commonly a mass spectrometer is combined with GC to generate a mass spectra which can be analysed or compared with a spectral database by computer for positive identification of each component in the mixture.


## Infrared spectroscopy

Certain bonds in a molecule absorb infra-red radiation at characteristic frequencies causing the covalent bonds to vibrate

Complicated spectra can be obtained than provide information about the types of bonds present in a molecule

ABOVE 1500  $\text{cm}^{-1}$  – “Functional group identification”

BELOW 1500  $\text{cm}^{-1}$  – “Fingerprinting”



Complicated and contains many signals – picking out functional group signals difficult.

This part of the spectrum is unique for every compound, and so can be used as a “fingerprint”.


e.g. C=O 1640 – 1750  $\text{cm}^{-1}$   
O-H (acid) 2500- 3300  $\text{cm}^{-1}$

A computer will compare the IR spectra against a database of known pure compounds to identify the compound


Use the IR absorption table provided in the data book to deduce presence or absence of particular bonds or functional groups

Bond	Wavenumber
C-O	1000-1300
C=O	1640-1750
C-H	2850 -3100
O-H Carboxylic acids	2500-3300 Very broad
N-H	3200-3500
O-H Acohols, phenols	3200- 3550 broad

use spectra to identify particular functional groups limited to data presented in wavenumber form e.g. an alcohol from an absorption peak of the O–H bond,



Spectra for butanal




Absorption or trough in between 1640-1750  $\text{cm}^{-1}$  range indicates presence of C=O bond


Always quote the wave number range from the data sheet

O-H absorptions tend to be broad

Absorption or trough in between 2500-3300  $\text{cm}^{-1}$  range indicates presence of O-H bond in an acid



Spectra for ethanoic acid




## Mass spectrometry

### Measuring the $M_r$ of an organic molecule

If a molecule is put through a mass spectrometer it will often break up and give a series of peaks caused by the fragments. The peak with the largest  $m/z$ , however, will be due to the complete molecule and will be equal to the  $M_r$  of the molecule. This **M** peak is called the **molecular ion**

### Spectra for $C_4H_{10}$



### M+1 peak

A small peak will often occur in a mass spectrum next to the molecular ion peak. This is due to the presence of the  $C^{13}$  isotope in the molecule.

The relative height of this **M+1** peak compared to the **M** peak can be used to calculate the number of carbons in a molecule. Use the following equation to calculate the number ( $n$ ) of carbons in a molecule

$$n = \frac{100 \times \text{height of M+1 peak}}{1.1 \times \text{height of M peak}}$$

Learn this equation

### Example

A hydrocarbon showed a ratio of heights of the **M** : **M+1** peaks of 9.3 : 0.5.

Calculate the number of carbon atoms present in one molecule of this hydrocarbon

$$n = \frac{100 \times \text{height of M+1 peak}}{1.1 \times \text{height of M peak}}$$

$$n = \frac{100 \times 0.5}{1.1 \times 9.3}$$

$$= 4.89$$

Round to 5 carbons

### M+2 peak

If a compound contains a chlorine or a bromine atom then a **M+2** peak will occur due to the two naturally occurring isotopes of chlorine or bromine.

Chlorine exists as  $Cl^{35}$  (75%) and  $Cl^{37}$  (25%)  
Bromine exists as  $Br^{79}$  (50%) and  $Br^{81}$  (50%)

$CH_3Cl$  will have a  $m/z$  value of **M** of 50  $CH_3Cl^{35}$  and **M+2** of 52  $CH_3Cl^{37}$   
The ratio of heights **M:M+2** will be 3:1

$CH_3Br$  will have  $m/z$  value of **M** of 94  $CH_3Br^{79}$  and **M+2** of 96  $CH_3Br^{81}$   
The ratio of heights **M:M+2** will be 1:1

If a compound contains two chlorine or bromine atoms then a **M+2** and a **M+4** peak will occur

$C_2H_4Cl_2$  will have a  $m/z$  value of **M** of 98  $C_2H_4Cl^{35}Cl^{35}$ , a **M+2** of 100  $C_2H_4Cl^{35}Cl^{37}$  and a **M+4** of 102  $C_2H_4Cl^{37}Cl^{37}$   
The ratio of heights **M:M+2: M+4** will be 9:6:1

$C_2H_4Br_2$  will have a  $m/z$  value of **M** of 186  $C_2H_4Br^{79}Br^{79}$ , a **M+2** of 188  $C_2H_4Br^{79}Br^{81}$  and a **M+4** of 190  $C_2H_4Br^{81}Br^{81}$   
The ratio of heights **M:M+2: M+4** will be 1:2:1

Relative abundances

$$Cl^{35}Cl^{35} = 0.75 \times 0.75 = 0.5625 \Rightarrow 9$$

$$Cl^{35}Cl^{37} + Cl^{37}Cl^{35} = 0.75 \times 0.25 \times 2 = 0.375 \Rightarrow 6$$

$$Cl^{37}Cl^{37} = 0.25 \times 0.25 = 0.0625 \Rightarrow 1$$

÷ smallest to get  
whole number ratio

$C_2H_3Cl_3$  will have a  $m/z$  value of **M** of 132  $C_2H_3Cl^{35}Cl^{35}Cl^{35}$ , a **M+2** of 134  $C_2H_3Cl^{35}Cl^{35}Cl^{37}$ , a **M+4** of 136  $C_2H_3Cl^{35}Cl^{37}Cl^{37}$  and a **M+6** of 138  $C_2H_3Cl^{37}Cl^{37}Cl^{37}$   
The ratio of heights **M:M+2:M+4:M+6** will be 27:27:9:1

Relative abundances

$$Cl^{35}Cl^{35}Cl^{35} = 0.75 \times 0.75 \times 0.75 = 0.4219 \Rightarrow 27$$

$$Cl^{35}Cl^{35}Cl^{37} + Cl^{35}Cl^{37}Cl^{35} = 0.75 \times 0.75 \times 0.25 \times 3 = 0.4219 \Rightarrow 27$$

$$Cl^{35}Cl^{37}Cl^{37} + Cl^{37}Cl^{35}Cl^{37} = 0.75 \times 0.25 \times 0.25 \times 3 = 0.1406 \Rightarrow 9$$

$$Cl^{37}Cl^{37}Cl^{37} = 0.25 \times 0.25 \times 0.25 = 0.0156 \Rightarrow 1$$

÷ smallest to get  
whole number ratio

## Fragmentation

When organic molecules are passed through a mass spectrometer, it detects both the whole molecule and fragments of the molecule.



The molecule loses an electron and becomes both an ion and a free radical


Several peaks in the mass spectrum occur due to fragmentation.

The Molecular ion fragments due to covalent bonds breaking:  $[M]^+ \rightarrow X^+ + Y\cdot$


This process produces an ion and a free radical. The ion is responsible for the peak

Relatively stable ions such as carbocations  $R^+$  such as  $\text{CH}_3\text{CH}_2^+$  and acylium ions  $[\text{R}-\text{C}=\text{O}]^+$  are common. The more stable the ion, the greater the peak intensity.



The peak with the highest mass/charge ratio will be normally due to the original molecule that hasn't fragmented (called the molecular ion). As the charge of the ion is +1 the mass/charge ratio is equal to Mr.




Equation for formation molecular ion







Equations for formation of fragment ions from molecular ions



Equation for formation molecular ion



Equations for formation of fragment ions from molecular ions



## NMR spectroscopy

### Different types of NMR

NMR spectroscopy involves interaction of materials with the low-energy radiowave region of the electromagnetic spectrum

NMR spectroscopy is the same technology as that used in 'magnetic resonance imaging' (MRI) to obtain diagnostic information about internal structures in body scanners

There are two main types of NMR


1.  $C^{13}$  NMR
2. H (proton) NMR

There is only around 1%  $C^{13}$  in organic molecules but modern NMR machines are sensitive enough to give a full spectra for  $C^{13}$   
The  $C^{13}$  spectra is a simpler spectrum than the H NMR


The radio waves used in proton nmr cause the hydrogen nucleus to **change its spin state.**

### Equivalent Carbon atoms.


In a  $C^{13}$  NMR spectrum, there is one signal (peak) for each **set of equivalent C atoms.**




3 peaks



4 peaks




4 peaks




2 peaks

1,2 dinitrobenzene




3 peaks

1,3 dinitrobenzene




4 peaks


1,4 dinitrobenzene



2 peaks



2 peaks










5 peaks

## Equivalent Hydrogen atoms.

In an H NMR spectrum, there is one signal for each set of equivalent H atoms.

In addition the **intensity (integration value)** of each signal is proportional to the **number of equivalent H atoms** it represents.



## Solvents

Samples are dissolved in solvents without any  $^1\text{H}$  atoms, e.g.  $\text{CCl}_4$ ,  $\text{CDCl}_3$ .

This means that in the H NMR the solvent will not give any peaks


The same solvent is used in  $\text{C}^{13}$  NMR and in this case there will be one peak due to the solvent that will appear on the spectrum. However, it is known where this peak is so it can be ignored. In the exam it is likely this peak will not occur on the spectra.

## Calibration and shift

A small amount of TMS (tetramethylsilane) is added to the sample to calibrate the spectrum

TMS is used because:


- its signal is away from all the others
- it only gives one signal
- it is non-toxic
- it is inert
- it has a low boiling point and so can be removed from sample easily



The same calibration compound is used for both H and C NMR





The spectra are recorded on a scale known as the chemical shift ( $\delta$ ), which is how much the field has shifted away from the field for TMS..

The  $\delta$  is a measure in parts per million (ppm) is a relative scale of how far the frequency of the proton signal has shifted away from that for TMS.



## H NMR shift

The shift depends on what other atoms/groups are near the H – more electronegative groups gives a greater shift.





## Proton exchange using D<sub>2</sub>O

If D<sub>2</sub>O is added to a sample then a process of proton exchange happens with the H in any O-H and N-H bonds. This has the effect of removing the peaks from the H-NMR spectra. This can help with the identification of O-H and N-H peaks on the spectra.


Addition of D<sub>2</sub>O to the sample of Propanoic acid would make the peak at  $\delta = 11.7$  (ppm) in the above spectrum disappear



### C<sup>13</sup> NMR shift



It will not be possible to identify the exact carbon corresponding to each peak if several carbons are in the same range








It is not possible to distinguish between similar shifts for each carbon in a benzene ring. In this example it should be possible to work out there are four different carbons in the benzene ring and these correspond to the four peaks between 120–145


## Spin-Spin coupling in H NMR

In high resolution H NMR each signal in the spectrum can be split into further lines due to inequivalent H's on neighbouring C atoms.

Splitting of peak = number of inequivalent H's on neighbouring C atoms + 1

signal	singlet	doublet	triplet	quartet	quintet
appearance					
Split number of peaks	1	2	3	4	5
number of neighbouring inequivalent H atoms	0	1	2	3	4
relative size		1:1	1:2:1	1:3:3:1	1:4:6:4:1

Nuclei in identical chemical environments do not show coupling amongst themselves!





The peak due to group **a** will be a **triplet** as it is next to **b** (a carbon with 2 H's)

The peak due to group **b** will be a **quartet** as it is next to **a** (a carbon with 3H's)

The peak due to group **c** will be a **singlet** as it is next to a carbon with no H's)

For 6 split peaks use the term hexet or multiplet



The peak due to group **a** will be a **singlet** as it is next to a carbon with 0 H's  
Shift 2.1-2.6  
Integration trace 3

The peak due to group **c** will be a **triplet** as it is next to a carbon with 2 H's  
Shift 0.7-1.2  
Integration trace 3


The peak due to group **b** will be a **quartet** as it is next to a carbon with 3 H's  
Shift 3.7 -4.1  
Integration trace 2

Hydrogens bonded to a Nitrogen or Oxygen usually do not couple with other protons and appear as singlets on the NMR spectra

The peak due to group **a** will be a **triplet** as it is next to a carbon with 2 H's  
Shift 0.7-1.2  
Integration trace 3

The peak due to group **b** will be a **quartet** as it is next to a carbon with 3 H's  
Shift 3.7 -4.1  
Integration trace 2

The peak due to group **c** will be a **singlet** as the Hydrogen is bonded to an oxygen and this does not split  
Shift 0.5-5.0  
Integration trace 1



You will not be asked to interpret splitting patterns for the protons attached to a benzene ring

## Bringing it all together

### 1. Work out empirical formula

Elemental analysis C 66.63% H 11.18% O 22.19%

C	H	O
66.63/12	11.18/1	22.19/16
=5.5525	=11.18	=1.386875
=4	=8	=1

### 2. Using molecular ion peak m/z value from mass spectrum calculate Molecular formula


molecular ion peak m/z value= 144

Mr empirical formula  $C_4H_8O = 72$

If Mr molecular formula 144 then compound is  $C_8H_{16}O_2$

### 3. Use IR spectra to identify main bonds/functional group


$C_8H_{16}O_2$  could be an ester, carboxylic acid or combination of alcohol and carbonyl. Look for IR spectra for C=O and O-H bonds



There is a C=O but no O-H absorptions, so must be an ester.


### 4. Use NMR spectra to give details of carbon chain

4 peaks – only 4 different environments.




Peak at  $\delta$  4 shows H-C-O

Area 2 suggests  $CH_2$   
Quartet means next to a  $CH_3$




Peak at  $\delta$  2.2 shows H-C=O

Area 2 suggests  $CH_2$   
Singlet means adjacent to C with no hydrogens



Peak at  $\delta$  1.2 shows R- $CH_3$   
Area 3 means  $CH_3$   
Triplet means next to a  $CH_2$



Put all together to give final structure

