2.4 Periodicity of Period 3

Trends in the reactions of the elements with water, limited to Na and Mg

Learn the equations

Sodium reacts with cold water. It fizzes around on surface etc.

2 Na (s) + 2 H₂O (l) \rightarrow 2 NaOH (aq) + H₂ (g)

Magnesium reacts very slowly with cold water to form the hydroxide but reacts more readily with **steam** to form the oxide Mg (s) + $H_2O(g) \rightarrow MgO(s) + H_2(g)$

Trends in the reactions of the elements Na, Mg, Al, Si, P and S with oxygen

The elements all react with oxygen to form oxides.

Sodium burns with a yellow flame to produce a white solid

Mg, Al, Si and P burn with a **white flame** to give **white solid** smoke

S burns with a **blue flame** to form an acidic **choking gas**.

4 Na (s) + O₂ (g) → 2 Na₂O (s) 2Mg (s) + O₂ (g) → 2MgO (s) 4AI + 3O₂ (g) → 2AI₂O₃ (s) Si + O₂ (g) → SiO₂ (s) 4P + 5O₂ (g) → P₄O₁₀ (s) S+ O₂ (g) → SO₂ (g) You should be able to write these equations.

Learn the formulae of the oxides

Sodium is stored under oil and phosphorus under water to stop these elements coming into contact and reacting with air.

A survey of the properties of the oxides of Period 3 elements

Understand the link between the physical properties of the highest oxides of the elements $Na \rightarrow S$ and their structure and bonding.

Ionic oxides

The metal oxides (Na₂O, MgO, Al₂O₃) are ionic. They have high melting points. They have **lonic giant lattice structures**: strong forces of attraction between ions: higher mp. They are ionic because of the large electronegativity difference between metal and O

The increased charge on the cation makes the ionic forces stronger (bigger lattice enthalpies of dissociation) going from Na to Al so leading to increasing melting points.

 ${\rm Al_2O_3}$ is ionic but does show some covalent character. This can be explained by the electronegativity difference being less big or alternatively by the small aluminium ion with a high charge being able to get close to the oxide ion and distorting the oxide charge cloud.

To prove that the above compounds contain ions experimentally - **melt** the solids and show they conduct electricity

Macromolecular oxides

SiO₂ is Macromolecular: It has many very strong covalent bonds between atoms. High energy needed to break the many strong covalent bonds – very high mp +bp

Simple molecular oxides:

 $P_4O_{10\;(s)},\ SO_{2\;(g)}$ are simple molecular with weak intermolecular forces between molecules (van der waals + permanent dipoles) so have lower mp's. They are covalent because of the small electronegativity difference between the non-metal and O atoms. P_4O_{10} is a molecule containing 4P's and 10 O's. As it is a bigger molecule and has more electrons than SO_2 it will have larger van der waals forces and a higher melting point.

Aluminium metal is protected from corrosion in moist air by a thin layer of aluminium oxide. The high lattice strength of aluminium oxide and its insolubility in water make this layer impermeable to air and water.

The reactions of the oxides of the elements Na -> S with water

Metal ionic oxides tend to react with water to form hydroxides which are alkaline

 $Na_2O(s) + H_2O(l) \rightarrow 2Na^+(aq) + 2OH^-(aq)$ pH 13 (This is a vigorous exothermic reaction)

The ionic oxides are basic as the oxide ions accept protons to become hydroxide ions in this reaction (acting as a bronsted

MgO (s) + H_2O (l) \rightarrow Mg(OH)₂ (s) **pH 9**

Mg(OH)₂ is only slightly soluble in water as its lattice is stronger so fewer free OH- ions are produced and so lower pH

Learn the equations!

Know the change in pH of the resulting solutions across the period.

Al₂O₃ and SiO₂ do not dissolve in water because of the high strength of the Al₂O₃ ionic lattice and the SiO₂ macromolecular structure, so they give a neutral pH 7

MgO is better than NaOH for treating acid in rivers and the stomach as it is only sparingly soluble and weakly alkaline so using an excess would not make the water excessively alkaline.

The non-metal, simple molecular, covalent, oxides react with water to give acids

 P_4O_{10} (s) + 6 H_2O (l) \rightarrow 4 H_3PO_4 (aq) pH 0 (this is a vigorous exothermic reaction) $SO_2 + H_2O \rightarrow H^+ + HSO_3^ SO_2(g) + H_2O(I) \rightarrow H_2SO_3(aq)$ pH 3 (weak acid)

 $SO_3(g) + H_2O(I) \rightarrow H_2SO_4(aq)$ 0 Ha

 $SO_3 + H_2O \rightarrow H^+ + HSO_4^-$

Equations showing formation of ions in solution

The trend is the **ionic metal oxides** show **basic** behaviour and the non-metal covalent oxides show acidic behaviour. The slightly intermediate nature of the bonding in aluminium oxide is reflected in its amphoteric behaviour: it can act as both a base and an acid

Displayed formulae of acids

Acid base reactions between period 3 oxides and simple acids and bases.

The basic oxides react with acids to make salts $Na_2O(s) + 2 HCI(aq) \rightarrow 2NaCI(aq) + H_2O(I)$ $Na_2O(s) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + H_2O(l)$ MgO (s) + 2 HCl (aq) \rightarrow MgCl₂ (aq) + H₂O (l) Or ionically

 $Na_2O(s) + 2H^+(aq) \rightarrow 2Na^+(aq) + H_2O(l)$

MgO (s) + 2 H⁺ (aq) \rightarrow Mg²⁺ (aq) + H₂O (l)

Rather than learning the equations by rote, learn the pattern. Most follow the pattern acid + base → salt + water Know the charges on the ions e.g. PO₄ 3- , SO₄2-

Amphoteric Oxides

Aluminium oxide can act as both an acid and an alkali and is therefore called amphoteric Aluminum oxide acting as a base

 Al_2O_3 (s)+ $3H_2SO_4$ (aq) $\rightarrow Al_2(SO_4)_3$ (aq) + $3H_2O$ (l)

 $Al_2O_3 + 6HCI \rightarrow 2AICl_3 + 3H_2O$

Or ionically $Al_2O_3 + 6H^+ \rightarrow 2Al^{3+} + 3H_2O$

Be careful for whether the question is asking for an ionic equation or a full one

Aluminum oxide acting as a acid

 Al_2O_3 (s)+ 2NaOH (ag) + $3H_2O$ (l) $\rightarrow 2NaAl(OH)_4$ (ag) (this equation needs learning carefully) Al_2O_3 (s)+ $2OH^-$ (aq) + $3H_2O$ (I) $\rightarrow 2Al(OH)_4$ (aq)

The other simple molecular acidic oxides react with bases to form salts.

 $P_4O_{10}(s) + 12 \text{ NaOH (aq)} \rightarrow 4Na_3PO_4(aq) + 6 H_2O(l)$ $P_4O_{10} + 6 Na_2O \rightarrow 4Na_3PO_4$

 $SO_2(g) + 2NaOH(aq) \rightarrow Na_2SO_3(aq) + H_2O(l)$

 $SO_3(g) + 2NaOH(aq) \rightarrow Na_2SO_4(aq) + H_2O(l)$

Or show ionically

 $P_4O_{10}(s) + 12 OH^-(aq) \rightarrow 4PO_4^{3-}(aq) + 6 H_2O(l)$

 $SO_2 (g) + 2OH^- (aq) \rightarrow SO_3^{2-} (aq) + H_2O (I)$

 $SO_3(g) + 2OH^-(aq) \rightarrow SO_4^{2-}(aq) + H_2O(l)$

SiO₂ has a giant covalent structure with very strong bonds. This stops SiO₂ dissolving or reacting with water and weak solutions of alkali. It will, however, react with very concentrated NaOH 2NaOH (I) + SiO₂ (s) \rightarrow Na₂SiO₃ (aq) + H₂O It is still classed as an acidic oxide